


RELATIVE RISK FOR TRANSMISSION FROM A PERSON LIVING WITH HIV¹



Decisions about sexual activity and condom use have a major effect on the risk for HIV transmission.

That's why it's so important to always practice safe sex.

This chart shows how the relative risk for a person living with HIV transmitting HIV to a person without the disease varies according to sexual activity and condom use. For example, insertive oral sex with a condom has a low risk for HIV transmission.

But receptive anal sex without a condom is 2,000 times riskier. Not having sex is the best way to protect against the transmission of HIV. But if you are having sex, it's important to know that all sex is not the same when it comes to transmitting HIV.^{1,2}



ANSWERS TO QUESTIONS YOUR PATIENTS MIGHT ASK³⁻⁶

Q: How does HIV cause AIDS?

A: HIV destroys certain kinds of blood cells (called CD4 cells) that are important to your immune system, which is what protects your body from disease. Without these cells, you become susceptible to many infections and some cancers.

Q: How is HIV spread?

A: The most common way that HIV is spread is sexual activity (anal, vaginal, or oral) with a person living with HIV. Sharing needles or works with a person living with HIV can also spread the disease. HIV can also be spread through transfusion of infected blood or blood products, though that risk is very low in the United States. Mothers with HIV can spread it to their babies before or during birth. HIV can also be spread through breast milk. Mothers with HIV should not breastfeed their babies.

Q: Can I pass HIV to someone else by doing things like shaking hands?

A: No. HIV is not spread through everyday contact at school, work, or in social settings. HIV cannot be caught from things like a handshake, a doorknob, a pet, or food. The virus does not live long outside the body.

Q: If I am living with HIV and pregnant, will my baby have HIV too?

A: Not necessarily. If you are treated with HIV drugs early in the pregnancy, the risk of passing the virus to your baby can be 2% or less. Even if you don't receive the HIV drugs until you are in labor, the chances of transmission are much lower than if you aren't treated at all. It's also important to remember that HIV can be spread through breast milk. Mothers with HIV should not breastfeed their babies.

Q: Is there a connection between HIV and other sexually transmitted diseases (STDs)?

A: Yes. If you become infected with another STD (such as syphilis, gonorrhea, or herpes), you are three to five times more likely to pass HIV to someone else than if you don't have another STD.

References:

1. Centers for Disease Control and Prevention. Incorporating HIV Prevention into the Medical Care of Persons Living with HIV: Recommendations of CDC, the Health Resources and Services Administration, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. *MMWR* 2003;52(RR-12):1–24.
2. Varghese B, Maher JE, Peterman TA, Branson, BM, Steketee RV. Reducing the risk of sexual HIV transmission: quantifying the per-act risk for HIV infection based on choice of partner, sex act, and condom use. *Sexually Transmitted Diseases* 2002;29:38–43.
3. Centers for Disease Control and Prevention. HIV/AIDS Questions and Answers. Available at: www.cdc.gov/hiv/resources/qa/index.htm. Accessed March 1, 2005.
4. Centers for Disease Control and Prevention. HIV testing among pregnant women—United States and Canada, 1998–2001. *MMWR* 2002;51(45):1013–1016.
5. Dorenbaum A, Cunningham CK, Gelber RD, et al. Two-dose intrapartum/newborn nevirapine and standard antiretroviral therapy to reduce perinatal HIV transmission: a randomized trial. *Journal of the American Medical Association* 2002;288:189–198.
6. Centers for Disease Control and Prevention. Is there a connection between HIV and other sexually transmitted diseases? Available at www.cdc.gov/hiv/pubs/faq/faq24.htm. Accessed March 9, 2005.